

Rabbit anti CD1a Polyclonal Antibody

Alternative Name(s): CD1; HTA1;

Order Information

• Description: CD1a • Catalogue: 500-4714 • Lot: See label • Size: 100ug/200ul • Host: Rabbit • Clone: nan • Application: IHC(P) • Reactivity: Hu

ANTIGEN PREPARATION

A synthetic peptide corresponding to internal sequence of human CD1a protein

BACKGROUND

CD1a is a 49 kD transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. CD1a plays a role in non-peptide glycolipid antigen presentation to CD1-restricted T cells. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. CD1a localizes to the plasma membrane and to recycling vesicles of the early endocytic system.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antbody recognizes human CD1a. The other species not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at –20oC to –70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 μg/ml
- ELISA: 0.01-0.1 μg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- · Flow cytometry: Not tested
- Molecular Weight: 47.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.



REFERENCES