

Rabbit anti LRRK2(pT1503) Polyclonal Antibody

Alternative Name(s): leucine-rich repeat kinase 2

Order Information

• Description: LRRK2(pT1503)

Catalogue: 630-060
Lot: See label
Size: 100ug/200ul
Host: Rabbit
Clone: nan

• Application: IHC(P), WB • Reactivity: Hu, Ms, Rt, Bv

ANTIGEN PREPARATION

leucine T1503)

BACKGROUND

LRRK2, leucine rich repeat kinase 2, belong to the leucine-rich repeat kinase family and encodes a protein with an ankryin repeat region, a leucine-rich repeat (LRR) domain, a kinase domain, a DFG-like motif, a RAS domain, a GTPase domain, a MLK-like domain, and a WD40 domain. The protein is present largely in the cytoplasm but also associates with the mitochondrial outer membrane. Mutations in this gene have been associated with Parkinson disease-8.

PURIFICATION

The Rabbit IgG is purified by site-modified Epitope Affinity Purification.

FORMULATION

This affinity purified antibody is supplied in sterile Tris-buffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human LRRK2(pT1503) protein with a phosphorylation site Threonine 1503. It cross reacts to human, mice and rat.

STORAGE


The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 μg/ml
- ELISA: 0.01-0.1 μg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- · Flow cytometry: Not tested
- Molecular Weight: 250.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.





Western Blot: The whole lysate derived from PMAstimulated U118MG (20 ug/lane) immunoblotted by Rabbit anti – LRRK2 (p1503) (Cat# 630-060) at 1:500. Observed a major immunoreactive band at molecular weight >200 kDa.

REFERENCES

FOR RESEARCH USE ONLY.