

AbboMax, Inc

Innovation at Work

Rabbit anti- GSK-3 β (CT) Polyclonal Antibody

Synonym: glycogen synthase kinase 3 beta (GSK3 β)

Order Information

Description: Rabbit anti -GSK-3beta (CT)
Catalogue#: 602-210
Lot#: See the label
Size: 100 ug/200 ul
Host: Rabbit
Clone: N/A
Application: ELISA, WB, IHC
Reactivity: Hu, Rt, Ms, Bv, Ck

ANTIGEN PREPARATION

A synthetic peptide corresponding to the C-terminus of GSK-3beta protein. This sequence is identical among human, mouse, rat, bovine, dog and chicken species.

BACKGROUND

Glycogen Synthase Kinase-3 β (GSK-3 β) is a serine/threonine kinase that affects glycogen metabolism by phosphorylating and down-regulating the activity of muscle glycogen synthase. Growth factor (such as Insulin) binds to its receptor, recruits PKB/AKT kinase to phosphorylate GSK-3beta on Serine 9, which prevents GSK-3beta from phosphorylating glycogen synthase. In contrast, the phosphorylation of tyrosine 216 in the catalytic domain increase this kinase activity to promote cell death under the absence of growth factors. GSK-3beta is also involved in Wnt signaling pathway where it forms a complex with axin, beta-catenin and adenomatous polyposis coli (APC) protein, thus the GSK-3beta inhibitor is considered as a potential approach in Alzheimer's Disease (AD) treatment.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification.

SPECIFICITY

This antibody recognizes GSK-3beta protein. It reacts with human, mouse and rat. The other species are not tested.

FORMULATION

This affinity purified antibody is supplied in sterile Phosphate-buffered saline (pH7.2) containing antibody stabilizer

STORAGE

The antibodies are stable for 12 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.


APPLICATIONS/SUGGESTED WORKING DILUTIONS

Western Blot	0.1-1 $\mu\text{g/ml}$
ELISA	0.01-0.1 $\mu\text{g/ml}$
Immunoprecipitation	2-5 $\mu\text{g/ml}$
IHC	Not tested
Flow cytometry	Not tested

MOLECULAR WEIGHT:	46 kDa
POSITIVE CONTROL:	Mouse muscle tissue lysate
CELLULAR LOCATION:	Cytoplasmic

Optimal dilutions should be determined by researchers for the specific applications.

DATA ATTACHMENTS



Western Blot: The mouse muscle tissue lysates were resolved onto 10% SDS-PAGE, transferred to NC membrane, and followed by an immunoblotting with Rabbit anti GSK-3beta (CT) (Cat#602-210) antibody (Lane 1) at 1:500, or with a pre-incubation of immunizing peptide (lane 2).

REFERENCES

Alan P. Kozikowski, et al. Highly Potent and Specific GSK-3 β Inhibitors That Block Tau Phosphorylation and Decrease α -Synuclein Protein Expression in a Cellular Model of Parkinson's Disease. ChemMedChem, 2006, 1(2), 256-266.

FOR RESEARCH USE ONLY.

AbboMax, Inc 1161 Ringwood Ct. Suite 100, San Jose, California 95131, USA
1 408-321-9898 (Tel). 1 408-321-9896 (Fax). 1-866-628-9898 www.abbomax.com info@abbomax.com