



Rabbit anti TGF-alpha Polyclonal Antibody

Alternative Name(s): transforming growth factor alpha

Order Information

- **Description:** TGF-alpha
- **Catalogue:** 500-3734
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), ELISA
- **Reactivity:** Hu

ANTIGEN PREPARATION

A synthetic peptide corresponding to the C-terminus of human TGF-alpha

BACKGROUND

Transforming growth factor alpha (TGF- α) is a member of the epidermal growth factor family. It is a ligand for the epidermal growth factor receptor, which activates a signaling pathway for cell proliferation, differentiation and development. TGF- α binding to EGFR promotes cell proliferation events that include wound healing and embryogenesis. TGF- α is also implicated in tumorigenesis and angiogenesis. It promotes tumorigenesis and regulates epithelial-mesenchymal transition modulation in colon cancer. TGF α has a higher expression in hepatocyte regeneration and proliferation in cirrhotic livers than in hepatocellular carcinoma.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human TGF-alpha . The other species not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 1900-01-05 12:00:00
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Immunohistochemistry: Human placenta (FFPE) stained with Rabbit anti-TGF-alpha (Cat#500-3734) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com