



Rabbit anti Actin, Smooth Muscle Polyclonal Antibody

Alternative Name(s): ACTA2

Order Information

- **Description:** Actin, Smooth Muscle
- **Catalogue:** 600-410
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu, Ms, Rt

ANTIGEN PREPARATION

A 10aa synthetic peptide corresponding to N-term of human Actin protein. This sequence is identical in human, mouse, rat, bovine and chicken

BACKGROUND

The Actin protein, a component of the eukaryotic cytoskeleton, is highly conserved between multiple species. It is involved in various types of cell motility, and maintenance of the cytoskeleton. In vertebrates, three main groups of actin isoforms, alpha, beta and gamma have been identified. The alpha actins are found in muscle tissues and are a major constituent of the contractile apparatus. The beta and gamma actins co-exist in most cell types as components of the cytoskeleton, and as mediators of internal cell motility. The levels of the Actin protein may vary between different cell types, but are consistent within a given cell type, and for this reason, Actin protein levels are considered an internal control for protein loading.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes actin protein. It cross-reacts to human, mouse, rat, chicken, bovine etc.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 42.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.


FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Western Blot: The whole lysate derived from Ht29 (20 ug/lane) immunoblotted by Rabbit anti -Actin (Cat# 600-410) at 1:500. Observed a major immunoreactive band at molecular weight ~42 kDa.



Immunohistochemistry: Human colon tissue (FFPE) stained with Rabbit Anti-Actin antibody (Cat# 600-410) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.