

Rabbit anti LRP6 (Non-phospho) Polyclonal Antibody

Alternative Name(s): Low density lipoprotein receptor-related protein 6

Order Information

• Description: LRP6 (Non-phospho)

Catalogue: 601-130
Lot: See label
Size: 100ug/200ul
Host: Rabbit
Clone: nan

Application: IHC(P), WB
 Reactivity: Hu/Rt/Ms

ANTIGEN PREPARATION

A synthetic peptide corresponding to the epitope KGTYF of human LRP6 protein

BACKGROUND

Low-desnsity lipoprotein receptor (LDLR)-related protein 6 (LRP6) is a member of the LDLR family which consists of cell surface proteins involved in receptor-mediated endocytosis of specific ligands. LRP5 cuases several human diseases including bone mineral density variability1, endosteal hyperostosis, autosomal dominant osteopetrosis type I etc. The V171 mutation in LRP5 causes high bone density by impairing the action of a normal antagonist of the Wnt pathway and thus increasing Wnt signaling. The activity of LRP5 and LRP6 can be inhibited by binding of some members of Dickkopf (DKK) family proteins. Upon stimulation with Wnt, LRP5 and LRP6 are phosphorylated at multiple sites including Thr 1479. Ser 1490 and Thr 1492 by kinases such as GSK-3 and CK1

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes ~180 kDa of human LRP6 protein. It also reacts with mouse and rat. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

• Western Blot: 0.1-1 µg/ml

• ELISA: 0.01-0.1 µg/ml

• Immunoprecipitation: 2-5 µg/ml

• IHC: 2-10 µg/ml

· Flow cytometry: Not tested


• Molecular Weight: 200.0

• Positive Control: Kidney Tissue

• Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.





Western Blot: The cell lysate derived from 3T3 was immunoprobed by Rabbit anti-LRP6 (Cat#601-130) at 1:500. An immunoreactive band is observed around ~180kDa.

REFERENCES

FOR RESEARCH USE ONLY.