

Rabbit anti H1N1(CT) (Influenza A virus) Polyclonal Antibody

Alternative Name(s): Haemagglutinin; HA; Influenza A virus; H1N1

Order Information

• Description: H1N1(CT) (Influenza A virus)

Catalogue: 601-240
Lot: See label
Size: 100ug/200ul
Host: Rabbit
Clone: nan

• Application: IHC(P), ELISA

• Reactivity: Virus

ANTIGEN PREPARATION

A synthetic peptide --PPLTPKQKREMAGTIRSV--derived from C-terminus of Influenza A virus nonstructural protein 1 (H1N1).

BACKGROUND

The Influenza A virus is a major public health threat. Novel influenza virus strains caused by genetic drift and viral recombination emerge periodically to which humans have little or no immunity, resulting in devastating pandemics. Influenza A can exist in a variety of animals; however it is in birds that all subtypes can be found. These subtypes are classified based on the combination of the virus coat glycoproteins hemagglutinin (HA) and neuraminidase (NA) subtypes. During 1997, an H1N1 avian influenza virus was discovered from Puerto Rico

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes H1N1(CT) (Influenza A virus) protein. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

• Western Blot: 0.1-1 μg/ml

• ELISA: 0.01-0.1 μg/ml

• Immunoprecipitation: 2-5 µg/ml

• IHC: 2-10 µg/ml

• Flow cytometry: Not tested


• Molecular Weight: 60.0

Positive Control: Kidney Tissue

• Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.





Western Blot: The recombinant protein derived from the full-length (225aa) of H5N1 (5 ug/lane) was resolved onto 10% of SDS-PAGE, transferred onto NC membrane, and immunoblotted by Rabbit anti –H1N1 (CT)(Cat#601-240) antibody at 1:500 . An immunoreactive band around ~30 kDa was observed.

REFERENCES