

Rabbit anti pTEN (Paired S380) Polyclonal Antibody

Alternative Name(s): Phosphatase and tensin homolog; PTEN; MMAC1

Order Information

• Description: pTEN (Paired S380)

Catalogue: 602-150
Lot: See label
Size: 100ug/200ul
Host: Rabbit
Clone: nan

Application: IHC(P), WB
 Reactivity: Hu, Ms, Rt

ANTIGEN PREPARATION

A synthetic peptide surrounding to the epitope -DTSDP—without a phosphorylation site at Ser385 of human PTEN protein. This sequence is identical among human, mouse, rat, bovine and chicken origins.

BACKGROUND

The phosphatase and tensin homologue (PTEN) tumor suppressor is a phosphatidylinositol D3-phosphatase that is frequently mutated in tumor cells. PTEN is itself regulated by phosphorylation on multiple serine and threonine residues in its C terminus. Casein kinase 2 (CK2) and Glycogen synthase kinase 3beta (GSK3beta), ATK/PI3K are involved in PTEN phosphorylation at sites of S370, S380, T382, T383, S385. The PTEN phosphorylation may play critical roles in the tumor suppressor functions.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes PTEN (Paired S385) without a phosphorylation site at Serine 385. It does not cross-react with phosphospecific peptides.

STORAGE


The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 μg/ml
- ELISA: 0.01-0.1 μg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- · Flow cytometry: Not tested
- Molecular Weight: 58.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

^{*}Optimal dilutions should be determined by researchers for the specific applications.





Western Blot: The mouse brain tissue lysate resolved onto 10% SDS-PAGE, transferred onto NC membrane, and followed by an immunoblotting with Rabbit anti PTEN (Paired S385) antibody (Cat#602-150) (Lane 1) at 1:500, or with a pre-incubation of immunizing peptide (lane 2). Observed a major immunoreactive band at molecular weight ~58kDa.

REFERENCES