



Rabbit anti FOXA3 Polyclonal Antibody

Alternative Name(s): Forkhead box A3

Order Information

- **Description:** FOXA3
- **Catalogue:** 602-700
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu, Ms, Rt

ANTIGEN PREPARATION

A synthetic peptide corresponding to the C-terminus of human FOXA3 protein. This sequence is identical to rat, mouse and human origins.

BACKGROUND

The forkhead box gene encodes a member of the forkhead class of DNA-binding proteins. These hepatocyte nuclear factors are transcriptional activators for liver-specific transcripts such as albumin and transthyretin, and they also interact with chromatin. Similar family members in mice have roles in the regulation of metabolism and in the differentiation of the pancreas and liver. FOXA1 plays an important role in breast cancer. The FOXA subfamily is composed of the transcription factors FOXA1, A2, A3. They were discovered as the regulators of hepatic genes. Foxa1 has been shown to be required for glucagon secretion in the pancreas, whereas Foxa2 is critical for the regulation of insulin secretion in pancreatic beta-cells. Foxa3 is important in the maintenance of glucose homeostasis, it is the dominating transcriptional regulator of GLUT2 expression in hepatocytes in vivo. In the normal activation of HNF-4alpha, HNF-1alpha, and PGC-1 induced by fasting is attenuated in mice lacking Foxa3.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes ~37 kDa of FOXA3 protein. It also reacts with human, mouse and rat. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 48.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com




*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



WB: Jurkat cell lysate
stained by RbxFOXA3
@ 1:2000 (Protein Amount:
lane 1: 20 ug lane 2: 10 ug
Lane3: 5 ug)

Western Blot: The whole lysate derived from HepG2 (20 ug/lane) immunoblotted by Rabbit anti – p53 (Cat#) at 1:500. Observed a major immunoreactive band at molecular weight ~48kDa.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com