



Mouse anti VEGFR2 (CD309) Monoclonal Antibody

Alternative Name(s): Vascular endothelial growth factor receptor 2 (VEGFR-2); KDR/ Flk-1; CD309 antigen

Order Information

- **Description:** CD309/VEGFR2
- **Catalogue:** 604-120
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Mouse
- **Clone:** 41Q23
- **Application:** IHC(P), WB
- **Reactivity:** Hu, Ms

ANTIGEN PREPARATION

A synthetic peptide from internal sequence of human CD309/VEGFR2 protein. This sequence is identical to mouse and rat.

BACKGROUND

Vascular endothelial growth factor receptor-2 (VEGFR-2), also known as CD309 antigen, the earliest known marker for vascular endothelia cells, is essential in vasculogenesis and angiogenesis. Binding of the dimeric VEGF-A to the extracellular domains of two monomeric receptors induces dimerization and activation of tyrosine kinase at tyrosine residues Y951, Y1054 and Y1059 in the kinase domain serve as positive regulatory sites. Phosphorylation at Y951 of the kinase insert is related to the migration of endothelial cells for tumor vascularization and growth. Cd309 is mainly expressed by endothelial cells, embryonic tissues, and megakaryocytes

PURIFICATION

The mouse IgG is purified by Protein A-Affinity Chromatography according to Isotyping

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human CD309/VEGFR2 protein. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*


- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: 149.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Immunohistochemistry: Human colon carcinoma (FFPE) stained with Mouse anti-CD309/VEGFR2 antibody (Cat# 604-120) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com