



Mouse anti CD99 Monoclonal Antibody

Alternative Name(s): MIC2; HBA71; MIC2X; MIC2Y; MSK5X

Order Information

- **Description:** CD99
- **Catalogue:** 605-090
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Mouse
- **Clone:** 3B2/TA8
- **Application:** IHC(P), FC
- **Reactivity:** Hu

ANTIGEN PREPARATION

A recombinant protein of human CD99

BACKGROUND

CD99 is a cell surface glycoprotein involved in leukocyte migration. It is type I single chain transmembrane protein devoid of N-linked glycosylation sites encoded by the pseudoautosomal gene MIC2. CD99 has an apparent molecular weight of 32 kD and is widely expressed on a variety of tissues. CD99 is highly expressed on thymocytes, T cells, and T cell leukemias and lymphomas. However, it is absent on some B cell lines, fetal B cells, eosinophils, granulocytes and the NK-cell line YT. CD99 is involved in spontaneous rosette formation with erythrocytes and may also be involved in other T-cell and hematopoietic cell adhesion pathways. CD99 has been reported to activate a caspase-independent death pathway in T cells under some conditions. CD99 interacts with a number of proteins including ferritin heavy chain 1, karyopherin beta 1, TRIP13, cyclophilin A, annexin II, and ubiquitin-conjugating enzyme E2H

PURIFICATION

The mouse IgG is purified by Protein A-Affinity Chromatography according to Isotyping

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human CD99 protein. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: 0.5-5 µg/10⁶ cells
- Molecular Weight: 40.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane


*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com



DATA ATTACHMENTS



Immunohistochemistry: Human Tonsil (FFPE) stained with Mouse anti- CD99 (Cat# 605-090) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com