



Mouse anti HLA-DR Monoclonal Antibody

Alternative Name(s): nan

Order Information

- **Description:** HLA-DR
- **Catalogue:** 605-481
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Mouse
- **Clone:** LN3
- **Application:** IHC(P), FC
- **Reactivity:** Hu

ANTIGEN PREPARATION

A recombinant protein of human HLA-DR

BACKGROUND

HLA-DR (Human Leukocyte Antigen – DR isotype) is $\alpha\beta$ heterodimeric cell surface glycoprotein, MHC class II cell surface receptor. Both α and β chains are anchored in the membrane, where α (heavy) chain is 36 kD and a β (light) chain is 27 kD. Each subunit contains two extracellular domains, a membrane-spanning domain and a cytoplasmic tail. The primary function of HLA-DR is to present peptide antigens. The human MHC class II molecules bind intracellularly processed peptides and present them to T-helper cells that lead to the production of antibodies against this peptide antigen. They therefore have a critical role in the initiation of the immune response. The complex of HLA-DR and peptide constitutes a ligand for the T-cell receptor (TCR). HLA-DR is expressed primarily on antigen presenting cells such as B lymphocytes, monocytes, macrophages, thymic epithelial cells and activated T lymphocytes. The reduced HLA-DR expression on monocytes is considered to correlate with infectious complications and the development of sepsis. Studies have shown that 3–20% of HLA-DR molecules constitutively reside in the lipid rafts of various antigen-presenting cells.

PURIFICATION

The mouse IgG is purified by Protein A-Affinity Chromatography according to Isotyping

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human HLA-DR protein. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 $\mu\text{g}/\text{ml}$
- ELISA: 0.01-0.1 $\mu\text{g}/\text{ml}$
- Immunoprecipitation: 2-5 $\mu\text{g}/\text{ml}$
- IHC: 2-10 $\mu\text{g}/\text{ml}$
- Flow cytometry: 0.5-5 $\mu\text{g}/10^6$ cells
- Molecular Weight: 34&28
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com




*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com



DATA ATTACHMENTS



Immunohistochemistry: Human Tonsil (FFPE) stained with Mouse anti-HLA-DR(Cat#605-481) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com