



Mouse anti beta-Catenin Monoclonal Antibody

Alternative Name(s): nan

Order Information

- **Description:** Catenin beta
- **Catalogue:** 605-540
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Mouse
- **Clone:** ZY480
- **Application:** IHC(P), WB
- **Reactivity:** Hu

ANTIGEN PREPARATION

A synthetic peptide of human beta-catenin

BACKGROUND

Beta-catenin is a cytosolic, 88 kDa, 781 amino acid protein belongs to the β -catenin family. β -catenin has been shown to perform two apparently unrelated functions: it has a crucial role in cell–cell adhesion in addition to a signaling role as a component of the Wnt/wg pathway. The N-terminus domain, containing the binding site and the phosphorylation sites. Beta-Catenin serves as a link between cytoskeleton actin and transmembrane cadherin(s). It can enter the nucleus and interact with the TCF/LEF family of transcription factors, initiating gene expression. Normally, β -catenin transcriptional activity is suppressed by a Ser/Thr kinase termed GSK3 β and/or Casein Kinase I (CK1). Kinases are constitutively active and phosphorylates β -catenin at multiple sites, including S33 and S37, Y96, Y228, Y280 etc. Phosphorylation of β -catenin targets the molecule for degradation via a ubiquitination-mediated pathway. GSK3 β activity can be blocked by upstream signaling events such as Wnt-Frizzled interaction. This inhibits GSK3 β , allowing unphosphorylated β -catenin to enter the nucleus and initiate gene activation. The phosphorylation of beta-catenin might contribute to tumorigenesis. Mutations in this gene are a cause of colorectal cancer (CRC), pilomatixoma (PTR), medulloblastoma (MDB), and ovarian cancer.

PURIFICATION

The mouse IgG is purified by Protein A-Affinity Chromatography according to Isotyping

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human Catenin beta protein. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 100.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com




*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com



DATA ATTACHMENTS



Immunohistochemistry: Human breast carcinoma (FFPE) stained with Mouse anti-catenin-beta (Cat# 605-540) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com