



## Mouse anti Tenascin C Polyclonal Antibody

Alternative Name(s): TNC; TN-C

### Order Information

- **Description:** Tenascin C
- **Catalogue:** 606-340
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Mouse
- **Clone:** T2H5
- **Application:** IHC(P), WB
- **Reactivity:** Ms, Rt, Hu

### **ANTIGEN PREPARATION**

A recombinant protein of human Tenascin C

### **BACKGROUND**

Tenascin-C (TN-C) is an extracellular matrix glycoprotein expressed during embryonic development, as well as in wound healing and cancer invasion in various tissues. It is usually derived from myofibroblasts in the cancer microenvironment. It has diverse functions, including weakening of cell adhesion, up-regulating the expression and activity of matrix metalloproteinases, modulating inflammatory responses, promoting recruitment of myofibroblasts, and enhancing fibrosis. TN-C could exert both harmful and protective effects and might be a therapeutic target as a key molecule in the control of the balance of beneficial and undesirable cellular responses during tissue remodeling. Cancer cell-specific TNC is a novel indicator of poor prognosis.

### **PURIFICATION**

The Rabbit IgG is purified by Epitope Affinity Purification

### **FORMULATION**

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

### **SPECIFICITY**

This antibody recognizes Tenascin C protein. It also cross reacts with human, mouse and rat tissues. The other species are not tested.

### **STORAGE**

The antibodies are stable for 24 months from date of receipt when stored at  $-20^{\circ}\text{C}$  to  $-70^{\circ}\text{C}$ . The antibodies can be stored at  $2^{\circ}\text{C}$ - $8^{\circ}\text{C}$  for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

### **APPLICATIONS/SUGGESTED WORKING DILUTIONS\***

- Western Blot: 0.1-1  $\mu\text{g/ml}$
- ELISA: 0.01-0.1  $\mu\text{g/ml}$
- Immunoprecipitation: 2-5  $\mu\text{g/ml}$
- IHC: 2-10  $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 250.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane


\*Optimal dilutions should be determined by researchers for the specific applications.

### **FOR RESEARCH USE ONLY.**

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA  
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). [www.abbomax.com](http://www.abbomax.com) [info@abbomax.com](mailto:info@abbomax.com)



## DATA ATTACHMENTS



Immunohistochemistry: Human brain (FFPE) stained with Mouse anti-Tenascin C (Cat# 606-340) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

## REFERENCES

### FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA  
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). [www.abbomax.com](http://www.abbomax.com) [info@abbomax.com](mailto:info@abbomax.com)