



Rabbit anti Tau Polyclonal Antibody

Alternative Name(s): microtubule-associated protein tau; MAPT

Order Information

- **Description:** Tau
- **Catalogue:** 620-790
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** hu, Ms, Rt

ANTIGEN PREPARATION

A synthetic peptide derived from C-terminus of human Tau protein. This sequence is identical to human, mouse, rat and bovine

BACKGROUND

Tau is a microtubule-associated phosphoprotein (MAP), localized in neuronal axons. It promotes tubulin polymerization and stabilizes microtubules. Tau proteins constitute a family of six isoforms which range from 352 to 441 amino acids. The tau variants differ from each other by the presence of either three or four repeat-regions in the carboxy-terminal part of the molecule and the absence or presence of one or two inserts in the amino-terminal part. Tau is hyperphosphorylated by ERK, GSK-3, TPKII and CDK5, at least thirty phosphorylation sites have been described, including Thr39, Ser46, Thr50, Thr69, Thr153, Thr175, Thr 181, Ser198, Ser199, Ser202, Thr205, Ser208, Ser210, Thr212, Ser214, Thr217, Thr231, Ser235, Ser237, Ser241, Ser262, Ser285, Ser305, Ser324, Ser352, Ser356, Ser396, Ser400, Thr403, Ser404, Ser409, Ser412, Ser413, Ser416 and Ser422. These sites are among the major abnormal phosphorylation sites of Tau. Phosphorylation on these sites reduces the ability of a given Tau species to promote microtubule self-assembly. Hyperphosphorylated Tau is the major protein of the paired helical filaments (PHFs), which make up the pathological neurofibrillary tangles of Alzheimer's disease (AD). The PHFs are also found in the lesions of other central nervous system disorders.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human Tau protein. It cross reacts to human, mouse and rat. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 µg/ml
- ELISA: 0.01-0.1 µg/ml
- Immunoprecipitation: 2-5 µg/ml
- IHC: 2-10 µg/ml
- Flow cytometry: Not tested
- Molecular Weight: ~46-80
- Positive Control: Kidney Tissue

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com




- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Western Blot: The tissue lysate derived from mouse brain was immunoblotted by Rabbit anti Tau (CT) (Cat#620-790) at 1:500. Immunoreactive bands are observed @ ~50-80 kDa.

REFERENCES

FOR RESEARCH USE ONLY.