



Rabbit anti AEG1(Paired S568) Polyclonal Antibody

Alternative Name(s): Metadherin; 3D3; AEG1; AEG-1; LYRIC; LYRIC/3D3

Order Information

- **Description:** AEG1(Paired S568)
- **Catalogue:** 630-090
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu, Ms, Rt, Bv

ANTIGEN PREPARATION

A synthetic peptide derived from a non-phosphorylated serine surrounding the epitope -TSWESPKQI-of human AEG1. This sequence is identical to human, mouse, rat and bovine.

BACKGROUND

AEG1 (Astrocyte elevated gene 1), also called MTDH (Metadherin), or L LYSine-Rich CEACAM1 co-isolated (LYRIC), is 582 aa single transmembrane domain (TMD) protein. It is widely expressed and highly conserved between species. It is recruited during the maturation of the tight junction complex and co-localizes with tight junction proteins ZO-1 and occludin in polarized epithelial cells. AEG-1 is involved in several crucial aspects of tumor progression, including transformation, evasion of apoptosis, invasion, metastasis, and chemoresistance. Overexpression of AEG-1 is frequently observed in melanoma, glioma, neuroblastoma, and carcinomas of breast, prostate, liver, and esophagus and is correlated with poor clinical outcomes. AEG-1 functions as a downstream mediator of the transforming activity of oncogenic Ha-Ras and c-Myc. Furthermore, AEG-1 overexpression activates the PI3K/Akt, nuclear factor κ B (NF κ B), and Wnt/ β -catenin signaling pathways to stimulate proliferation, invasion, cell survival, and chemoresistance. The lung-homing domain of AEG-1 also mediates the adhesion of tumor cells to the vasculature of distant organs and promotes metastasis. These findings suggest that therapeutic targeting of AEG-1 may simultaneously suppress tumor growth, block metastasis, and enhance the efficacy of chemotherapeutic treatments.

PURIFICATION

The Rabbit IgG is purified by Epitope Affinity Purification

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes Non-phosphospecific AEG1 at surrounding Serine 568. It does not recognize phosphospecific site pSer568. This antibody reacts with human, mouse and rat origins. The other species are not tested.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 64.0

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com



- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane

*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com

DATA ATTACHMENTS



Western Blot: The cell lysate derived from HELA cells was immuno-blotted by Rabbit anti Non-phosphospecific AEG1 (Paired S568) (Cat#630-090) at 1:500. An immunoreactive band around 64kDa is observed.

REFERENCES

FOR RESEARCH USE ONLY.