



Rabbit anti STK17b Polyclonal Antibody

Alternative Name(s): STK17b, Drak2

Order Information

- **Description:** STK17b
- **Catalogue:** 630-360
- **Lot:** See label
- **Size:** 100ug/200ul
- **Host:** Rabbit
- **Clone:** nan
- **Application:** IHC(P), WB
- **Reactivity:** Hu, Ms, Rt

ANTIGEN PREPARATION

A synthetic peptides from N-terminus of human STK17b. This sequence is identical among human, rat and mouse.

BACKGROUND

STK17b, a serine/threonine kinase 17b, is also called DRAK2. DRAK2 serves as a TGF- β 1-inducible antagonist of TGF- β signaling. TGF- β 1 stimulation rapidly induces DRAK2 expression and enhances endogenous interaction of the type I TGF- β receptor with DRAK2, thereby blocking R-Smads recruitment. Depletion of DRAK2 expression markedly augmented the intensity and the extent of TGF- β 1 responses. Furthermore, a high level of DRAK2 expression was observed in basal-like and HER2-enriched breast tumors and cell lines, and depletion of DRAK2 expression suppressed the tumorigenic ability of breast cancer cells. DRAK2 is an intrinsic intracellular antagonist participating in the negative feedback loop to control TGF- β 1 responses, and aberrant expression of DRAK2 increases tumorigenic potential, in part, through the inhibition of TGF- β 1 tumor suppressor activity.

PURIFICATION

The Rabbit IgG is purified by site-modified Epitope Affinity Purification.

FORMULATION

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

SPECIFICITY

This antibody recognizes human STK17b protein. It cross reacts to human, mice and rat.

STORAGE

The antibodies are stable for 24 months from date of receipt when stored at -20°C to -70°C . The antibodies can be stored at 2°C - 8°C for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

APPLICATIONS/SUGGESTED WORKING DILUTIONS*

- Western Blot: 0.1-1 $\mu\text{g/ml}$
- ELISA: 0.01-0.1 $\mu\text{g/ml}$
- Immunoprecipitation: 2-5 $\mu\text{g/ml}$
- IHC: 2-10 $\mu\text{g/ml}$
- Flow cytometry: Not tested
- Molecular Weight: 69.0
- Positive Control: Kidney Tissue
- Cellular Location: Cell Membrane


*Optimal dilutions should be determined by researchers for the specific applications.

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com



DATA ATTACHMENTS



Immunohistochemistry: The whole cell pellet MOLT4 (FFPE) stained with Rabbit anti-STK17b antibody (Cat# 630-360) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.

REFERENCES

FOR RESEARCH USE ONLY.

AbboMax, Inc 2528 Qume Drive, Suite 8, San Jose, California 95131, USA
1 408-573-1898 (Tel). 1 408-573-1858 (Fax). www.abbomax.com info@abbomax.com