

# Rabbit anti MSH6 Polyclonal antibody

Alternative Name(s): nan

#### **Order Information**

Description: MSH6
Catalogue: 630-850
Lot: See label
Size: 100ug/200ul
Host: Rabbit
Clone: nan

Application: IHC(P), WB
Reactivity: Hu, Ms, Rt,

# **ANTIGEN PREPARATION**

A synthetic peptide from internal sequence of human MSH6. it is identical to human, mice and rat.

#### **BACKGROUND**

MSH6 is a member of the DNA mismatch repair MutS family. Defects in MSH6 are a cause of hereditary non-polyposis colorectal cancer (HNPCC) (Lynch syndrome). HNPCC is an autosomal, dominantly inherited disease associated with marked increase in cancer susceptibility. It is characterized by a familial predisposition to early onset colorectal carcinoma (crc) and extra-colonic cancers of the gastrointestinal, urological and female reproductive tracts. HNPCC is reported to be the most common form of inherited colorectal cancer in the western world. MSH6 is central to mismatch DNA repair.

# **PURIFICATION**

The Rabbit IgG is purified by Epitope Affinity Purification

#### **FORMULATION**

This affinity purified antibody is supplied in sterile Phosphatebuffered saline (pH7.2) containing antibody stabilizer

#### SPECIFICITY

This antibody recognizes human MSH6 protein. It cross reacts to human, mice and rat.

# **STORAGE**

The antibodies are stable for 24 months from date of receipt when stored at -20oC to -70oC. The antibodies can be stored at 2oC-8oC for three month without detectable loss of activity. Avoid repeated freezing-thawing cycles.

### APPLICATIONS/SUGGESTED WORKING DILUTIONS\*

• Western Blot: 0.1-1 μg/ml

• ELISA: 0.01-0.1 μg/ml

• Immunoprecipitation: 2-5 µg/ml

• IHC: 2-10 µg/ml

· Flow cytometry: Not tested


• Molecular Weight: 153.0

Positive Control: Kidney Tissue

• Cellular Location: Cell Membrane

<sup>\*</sup>Optimal dilutions should be determined by researchers for the specific applications.





Immunohistochemistry: Human placenta (FFPE) stained with Rabbit anti-MSH6 antibody (Cat# 630-850) at 1:200 for 10 min @ RT. Staining of formalin-fixed tissue requires boiling tissue sections in 10 mM Citrate Buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min

# **REFERENCES**